

THE UROLOGY GROUP

www.urologygroupvirginia.com

1860 Town Center Drive · Suite 150/160 · Reston, VA 20190 · 703-480-0220
19415 Deerfield Avenue · Suite 112 · Leesburg, VA 20176 · 703-724-1195
224-D Cornwall Street, NW · Suite 400 · Leesburg, VA 20176 · 703-443-6733
24430 Stone Springs Blvd · Ste 100 · Dulles · VA 20166

PARTIAL NEPHRECTOMY POSTOPERATIVE INSTRUCTIONS

Hospital stay: Patients typically stay overnight in the hospital for 1-2 days for monitoring, intravenous fluids and pain management.

Diet: After anesthesia, begin with clear liquids. You may take what you like to eat or drink. Depending on how you feel the following day, you may resume your normal diet. The appetite may be diminished the first several days at home. Drink plenty of water and avoid heavy meals.

Activity: *Be sure to walk at least three times per day.* This helps prevent blood clots in the legs, which can travel to the lung and become life-threatening. You may take walks outside. You may go up and down stairs. Your energy will return over the course of four to six weeks.

You should avoid strenuous activity for 6-8 weeks. This includes activities such as golf, tennis, cutting the grass, etc. You should avoid carrying anything over fifteen pounds for 6-8 weeks.

Surgical drain: A surgical drain called a Jackson Pratt or JP drain will be in place after surgery to drain excess fluid and blood. This will be removed before leaving the hospital or at your postop appointment.

Bathing: You may take a shower as you normally do. You may allow the water to run over the incision then pat dry. Avoid submerging in water for two days. You may have staples in the incision. These will be removed postoperatively. You do not need to place anything over the incision but may place a gauze if there is oozing or spotting.

Medication: You can obtain good pain relief by taking two acetaminophen (Tylenol) every four hours while awake for the first several days. You will also get a prescription for pain pills. You can use these prescription pain pills in addition to acetaminophen every four hours. Do not exceed 4000 mg acetaminophen per day.

You may receive a prescription for a stool softener to avoid straining after surgery. Take plenty of fiber and water or over the counter stool softener (Senekot, Milk of Magnesia) to avoid straining for bowel movements. It may take a few days to have your first bowel movement after surgery.

Postoperative appointment: You will need a postoperative visit in approximately 7-10 days after your discharge. Surgical staples and drain may be removed at this time. Call the office to make an appointment if you do not already have one.

Call the office or come to the emergency room for fever over 101°F, bright red blood in the urine, difficulty breathing, chest pain, dizziness, nausea, vomiting, leg swelling or pain.